



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NUMBER THEORY.

215. Proposed by R. D. CARMICHAEL, Indiana University.

Find one or more values of n such that a polygon of n sides shall have the number of its diagonals equal to the cube of an integer.

SOLUTION BY WALTER C. EELLS, U. S. Naval Academy.

Since the number of diagonals of an n -sided polygon is $\frac{n(n-3)}{2}$, this is equivalent to the problem: find solutions in integers of $n^2 - 3n = 2k^3$. By the aid of a table of squares it is easily found that the only values of n less than 1,000 satisfying this equation are 9 and 128, for which $k = 3$ and 20 respectively, the number of diagonals being 27 and 8,000.

216. Proposed by ELIJAH SWIFT, University of Vermont.

If p is a prime > 3 , show that $\sum_{a=1}^{a=p-1} 1/a \equiv 0 \pmod{p^2}$, where $1/a \equiv x$, if $ax \equiv 1 \pmod{p^2}$.

SOLUTION BY THE PROPOSER.

Referring to my solution¹ of algebra problem number 385, I proved that

$$A_{p-2} \equiv 0 \pmod{p^2}, \text{ where } A_{p-2} \equiv 1 \cdot 2 \cdot 3 \cdots p-2 + 1 \cdot 3 \cdot 4 \cdots p-1 + \cdots$$

Hence,

$$A_{p-2} \equiv (1 \cdot 2 \cdot 3 \cdots (p-1)) \sum_{a=1}^{p-1} \frac{1}{a}.$$

Suppose that

$$1 \cdot 2 \cdot 3 \cdots (p-1) \equiv -1 + A \cdot p \pmod{p^2}.$$

Then

$$A_{p-2} \equiv \sum_{a=1}^{p-1} \frac{-1 + Ap}{a} = \sum_{a=1}^{p-1} \frac{-1}{a} + pA \sum_{a=1}^{p-1} \frac{1}{a}.$$

But

$$\sum_{a=1}^{a=p-1} \frac{1}{a} \equiv \sum_{a=1}^{a=p-1} a \pmod{p} \equiv 0,$$

since

$$A_{p-2} \equiv 0 \pmod{p^2} \quad \sum_{a=1}^{a=p-1} \frac{1}{a} \equiv 0 \pmod{p^2}.$$

NOTES AND NEWS.

EDITED BY W. DEW. CAIRNS.

MISS MARIE GUGEL, formerly teacher of mathematics in the Toledo, Ohio, high school, is now supervisor of high schools in Columbus. She is secretary of the mathematics section of the Central Association of Science and Mathematics Teachers.

MR. FORREST R. BAKER, assistant in mathematics at the University of Michigan, died December 6, following an operation for appendicitis.

¹ Volume XXI, page 157, May, 1914.